




Daran
on RYTARY
since 2015



Denise
on RYTARY
since 2016



David
on RYTARY
since 2016

**COME
JOIN US!**

Moving ON With RYTARY

A Community-Based Conversation

Join us for an interactive conversation between a Stories in Parkinson's (SPARK) Ambassador and a healthcare provider! Together, they will share insights about living well with Parkinson's, managing symptoms, and how RYTARY might be able to help. But that's not all. At this event, you will:



Hear firsthand from a SPARK Ambassador about their experience with Parkinson's and where they find inspiration



Explore an exciting Parkinson's treatment option and resources for talking to your healthcare provider



Connect with others in the Parkinson's community and find answers to your questions

Who is invited?

People living with Parkinson's disease and their care partners

When is it?

*Please plan to log in 15 minutes early

Where will it

*Please click the link above to join the webinar

Who will present?

accompanied by a SPARK Ambassador

**SIGN
UP NOW!**

Visit takingonpd.com or Call **1-844-947-4211**

*Thank you for your interest!
We look forward to welcoming you at our program!*

Sponsored by Amneal

INDICATION

RYTARY (carbidopa and levodopa) extended-release capsules is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

Do not take RYTARY with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors because taking these two drugs within two weeks of each other can result in high blood pressure.

Please see additional Important Safety Information on back and [full Prescribing Information](#) on RYTARY.com

RYTARY[®]
(carbidopa and levodopa)
EXTENDED-RELEASE CAPSULES
23.75 mg/95 mg • 36.25 mg/145 mg
48.75 mg/195 mg • 61.25 mg/245 mg

IMPORTANT SAFETY INFORMATION (continued)

Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting RYTARY. This may affect your ability to drive or operate machinery. Do not do anything that requires alertness until you know how RYTARY affects you.

Tell your healthcare provider if you have any heart conditions, especially if you have had a heart attack or irregular heartbeats; if you experience hallucinations or abnormal thoughts and behaviors (such as excessive suspicion, believing things that are not real, confusion, agitation, aggressive behavior, and disorganized thinking), if you have intense urges to gamble, increased sexual urges, other intense urges, and the inability to control those urges; if abnormal involuntary movements appear or get worse during treatment with RYTARY; or if you have ever had a peptic ulcer or glaucoma.

The most common side effects that may occur with RYTARY include nausea, dizziness, headache, sleeplessness, abnormal dreams, dry mouth, abnormal involuntary movements, anxiety, constipation, vomiting, and low blood pressure upon rising.

Some patients taking RYTARY have experienced suicidal thoughts or have attempted suicide. Tell your healthcare provider if you have thoughts of suicide or have attempted suicide.

Notify your healthcare provider if you become pregnant or intend to become pregnant during therapy or if you intend to breast-feed or are breast-feeding an infant.

Talk to your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle stiffness.

Take RYTARY as instructed. You may take RYTARY with or without food; however, taking RYTARY with food may decrease or delay its effect. For this reason, consider taking the first dose of the day about 1 to 2 hours before eating. Swallow RYTARY whole; do not chew, divide, or crush. If you have difficulty swallowing the capsule, twist apart both halves and sprinkle the entire contents of both halves of the capsule on a small amount of applesauce (1 to 2 tablespoons). Consume the mixture immediately. Do not store the drug/food mixture for future use.

Note: This information is intended to aid in the safe and effective use of RYTARY. It is not a disclosure of all possible adverse or intended effects. Tell your healthcare provider if you have any side effects while taking RYTARY. He or she can make adjustments that may reduce these effects.

To report SUSPECTED ADVERSE REACTIONS, contact Amneal Specialty, a division of Amneal Pharmaceuticals LLC at 1-877-835-5472 or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please read the full Prescribing Information. For more information go to RYTARY.com and/or talk to your healthcare provider.



RYTARY® is a registered trademark of Amneal Pharmaceuticals LLC.
© 2021 Amneal Pharmaceuticals LLC. Distributed by Amneal Specialty,
a division of Amneal Pharmaceuticals LLC.
All rights reserved. PP-SPM-RYT-US-0164 03/2021

RYTARY[®]
(carbidopa and levodopa)
EXTENDED-RELEASE CAPSULES
23.75 mg/95 mg • 36.25 mg/145 mg
48.75 mg/195 mg • 61.25 mg/245 mg